**Supplement 1-Addendum:** Appendix (Questionnaire)

## General section

### 1. Gender

a) Male

b) Female

### 2. Age

a) 18-25 years

b) 26-35 years

c) 36-45 years

d) 46-55 years

e) 56-65 years

f) 66-75 years

g) 76 years and older

## Part I - Usual activities

3. a) I have no problems performing everyday activities.

b) I have minor problems performing everyday activities.

c) Performing everyday activities is a moderate problem for me.

d) Performing everyday activities is a major problem for me.

e) I am unable to perform everyday activities.

4. a) I have no problems walking or strolling.

b) Walking or strolling is a minor problem for me.

c) Walking or strolling is a moderate problem for me.

d) Walking or strolling is a major problem for me.

e) I am unable to walk.

## Part II - Pain/discomfort

5. a) I have no pain or discomfort in my chest.

   b) I have mild chest pain or discomfort.

   c) I have moderate pain or discomfort in the chest.

   d) I have severe pain or discomfort in the chest.

   e) I have very severe pain in my chest.

## Part III - Anxiety and Depression


6. a) I am not anxious or depressed.

   b) I am mildly anxious or depressed.

   c) I am moderately anxious or depressed.

   d) I am severly anxious or depressed

   e) I am very severely anxious or depressed




VAS scale The VAS scale is used to assess the intensity of symptoms and the QoL of patients. Patients mark their subjective experience on a 10 cm long line, where the endpoints represent the extreme values of the symptoms.